

Data Standard	Single-Pool Urea Kt/V (spKt/V) and Equilibrated Urea Kt/V (eKt/V) Dialysis Adequacy Measures for a Single Hemodialysis Treatment
Alternate Name(s)	Single-Pool Urea Kt/V for Hemodialysis spKt/V Method: Second Generation Ultrafiltration-Adjusted Estimation eKt/V Method: Patient Clearance Time Correction
Description	<p>The single-pool urea Kt/V for hemodialysis is a dimensionless measure of the adequacy of small molecule removal provided by a single dialysis treatment, where K is the dialyzer urea clearance, t is the treatment time, and V is the urea distribution volume for the patient.</p> <p>As discussed in the Rationale section below, spKt/V can be estimated from 5 parameters from a single hemodialysis treatment. This formula represents one such calculation, based on a second-generation ultrafiltration-adjusted estimate, which is commonly used today.</p> <p>Two related hemodialysis adequacy measures are derived from the spKt/V:</p> <ul style="list-style-type: none"> • Equilibrated Kt/V or double-pool Kt/V, which accounts for the rebound in the blood urea nitrogen concentration after completion of the dialysis treatment. • Standard weekly Kt/V, which is a measure of total clearance per week, and as such, accounts for the number of treatments during the week <p>Separate specifications and appropriate applications for those metrics are provided below.</p>
Rationale	<p>An easily measured product of protein breakdown, urea has been an accepted marker for determination of small-molecule clearance in dialysis for over thirty years.¹⁻⁶ The Kt/V term arises from solution of a very simple time-dependent mass balance on urea nitrogen based on the following assumptions:</p> <ul style="list-style-type: none"> • The human body acts as a single well-mixed reservoir in which urea is uniformly distributed. • The volume of that reservoir is constant during the dialysis treatment (i.e., ignore ultrafiltration) • The rate of change of urea concentration with time is proportional to the urea concentration. <p>The urea distribution volume of a patient depends on body size, gender, and body composition. It is costly and impractical to measure each patient's urea</p>

	<p>distribution volume by indicator dilution. However, if the dialyzer clearance is known, then V can be calculated from the pre/post BUN, pre/post weight, and treatment time by iterative solution using formal urea kinetic modeling (UKM). UKM relies on an iterative method to determine the urea distribution volume.</p> <p>Given the challenges related to such an iterative process, various empirically derived equations have been developed for estimating spKt/V. In recent years, the so-called “Daugirdas II” Kt/V approximation has been in wide use. For example, the equation is one of the accepted dialysis dose methods acceptable to the U.S. Centers for Medicare and Medicaid Services (CMS) for claims and other reporting. First published in 1989, the original Daugirdas formula for estimating variable-volume spKt/V for a dialysis treatment frequency of 3x per week was found to overestimate Kt/V for values of Kt/V above.^{1,3,4} The revised equation from 1993 provided more accurate Kt/V estimates at higher Kt/V’s.⁵ The following specifications are based on the second-generation equation.</p>
Data Source(s)	<ul style="list-style-type: none"> • Hemodialysis treatment-level data • Laboratory results
Required Data Elements	<ul style="list-style-type: none"> • Pre-dialysis weight (PreWt, kg) • Post-dialysis weight (PostWt, kg). The weighing conditions should be comparable for the pre- and post-dialysis weight (e.g., the same scale, identical clothing worn, use of a wheelchair, etc.) • Pre-dialysis Blood Urea Nitrogen (PreBUN) concentration (mg/dL) • Post-dialysis Blood Urea Nitrogen (PostBUN) concentration (mg/dL). For accurate results, the “Stop Flow/Stop Pump” technique should be used, in which the blood flow rate is stopped prior to drawing blood for the PostBUN sample. • Delivered treatment time (t, minutes). The delivered treatment time reflects the total time of administered dialysis (i.e., excludes time for intermittent dialysis discontinuation if blood returned by setting the machine in bypass mode). It should also exclude any time during the procedure when the patient is not connected to the machine and receiving dialysis. For example, the treatment time should be adjusted if dialysis is temporarily interrupted (e.g., to allow the patient to use bathroom facilities).
Calculation Method	<p>Single Treatment spKt/V for HD using 2nd generation UFR-adjusted estimate⁵</p> $\text{spKt/V} = -\ln \left[\frac{\text{PostBUN}}{\text{PreBUN}} - (0.008 \times t/60) \right] + [4 - (3.5 \times \text{PostBUN}/\text{PreBUN})] \times [(\text{PostWt} - \text{PreWt})/\text{PostWt}]$ <p>Single Treatment eKt/V for HD using patient clearance time concept⁷ with modified clearance time constant⁸</p> $\text{eKt/V} = \text{spKt/V} \times t/(t + 30.7)$

	<p>Alternative Single Treatment eKt/V for HD using multiple linear regression⁶</p> $eKt/V = spKt/V * [0.924 - 0.395 / (t /60)] + 0.056$ <p>Both spKt/V and eKt/V for HD are typically rounded to the first decimal point (e.g., 1.34 rounds to 1.3 and 1.35 rounds to 1.4) since the BUNs are usually recorded with two significant digits (e.g., 65 mg/dL and 21 mg/dL). However, unrounded spKt/V values should be used in eKt/V calculations.</p>
<p>Exclusions</p>	<ul style="list-style-type: none"> • Hemodialysis treatments without the required data elements • Pediatric patients.
<p>Additional Desirable Data Elements for Collection</p>	<ul style="list-style-type: none"> • Method of Kt/V calculation • Specific hemodialysis modality: in-center hemodialysis, home hemodialysis, home nocturnal hemodialysis, or in-center nocturnal hemodialysis (i.e., >6-hour treatment time in-center).
<p>Notes</p>	<ul style="list-style-type: none"> • If a researcher chooses to aggregate and calculate the average spKt/V HD for any time period (e.g., 30-days, 3 months, etc.), first calculate the spKt/V for each individual treatment and then calculate the average by summing the spKt/V's across treatments and dividing by the total number of treatments. • Where possible, conduct data cleaning of input values prior to Kt/V calculation. For example, remove atypically low PostBUN values that may represent blood draw errors. If that is not possible, consider removing atypically high Kt/V values and exercise particular caution when averaging values over long periods of time. • For the purposes of research, each individual data element and the calculated spKt/V HD should be collected. This allows identification of outlier input parameters such as a diluted post BUN or a weight change during dialysis that exceeds a clinically reasonable range. • Single pool Kt/V is generally used for thrice weekly hemodialysis whereas standardized Kt/V can be used to estimate clearance for any frequency of dialysis.
<p>Example Measure Calculation</p>	<p>The following is an example of how to calculate spKt/V for HD using the above equation:</p> <p>In-center hemodialysis modality, prescribed thrice weekly dialysis Pre-dialysis weight 73 kg, post-dialysis weight 70 kg, total treatment time 240 minutes Pre-dialysis BUN 60 mg/dL, Post-dialysis BUN 18 mg/dL</p>

	$\text{spKt/V} = -\ln(18/80 - 0.008 \cdot 240/60) + ((4 - 3.5 \cdot (18/80)) \cdot (3/70)) = 1.443$, which rounds to 1.4 $\text{eKt/V} = 1.443 \times (240)/(240 + 30.7) = 1.279$, which rounds to 1.3 or alternatively: $\text{eKt/V} = 0.924 \cdot 1.443 - 0.395 \cdot 1.443/(240/60) + 0.056 = 1.247$, which rounds to 1.2
Acronyms	spKt/V: Single pool Kt/V eKt/V: Equilibrated Kt/V HD: Hemodialysis
Synonyms	Daugirdas II Kt/V is used interchangeably with single pool Kt/V

References

1. Wineman RJ: Artificial kidney-chronic uremia program: Plans for cooperative clinical trials. *Kidney Int* 1975.7(Suppl 2):S243-S246,.
2. Laird NM, Berkey CS, Lowrie EG: Modeling success or failure of dialysis therapy: The National Cooperative Dialysis Study. *Kidney Int* 1983.23(Suppl 13):S101-S107.
3. Gotch FA and Sargent JA: A mechanistic analysis of the National Cooperative Dialysis Study (NCDS). *Kidney Int* 1985.28: 526-534.
4. Daugirdas JT: The post: pre-dialysis plasma urea nitrogen ratio to estimate K.t/V and NPCR: mathematical modeling. *Int J Artif Organs*.1989. Jul;12(7):411-9.
5. Daugirdas JT: Second generation logarithmic estimates of single pool variable volume Kt/V: an analysis of error. *J Am Soc Nephrol* 1993.4(5) Nov:1205-13.
6. Leypoldt JK, Jaber BL, and Zimmerman, DL. Predicting Treatment Dose for Novel Therapies Using Urea Standard Kt/V. *Sem Dial* 2004.17(2):142-145.
7. Tattersall JE, DeTakats D, Chamney P, Greenwood RN and Farrington K. The post-hemodialysis rebound: Predicting and quantifying its effect on Kt/V. *Kidney Int* 1996. 50:2094-2102.
8. Greene T, Daugirdas JT, Depner TA, Gotch F, Kuhlman M. Frequent Hemodialysis Network Study Group: Solute clearances and fluid removal in the frequent hemodialysis network trials. *Am J Kidney Dis* 2009.53: 835-844.